

# Understanding Quantum Information and Computation

Lesson 9

## Density Matrices

---

John Watrous

# Motivation

- Density matrices represent a *broader class* of quantum states than quantum state vectors.
- Density matrices can describe states of *isolated parts* of systems, such as the state of one system that happens to be entangled with another system that we wish to ignore.
- Probabilistic states can be represented by density matrices, allowing quantum and classical information to be described together within a *single mathematical framework*.

# Definition of density matrices

Suppose that  $X$  is a system and  $\Sigma$  is its classical state set.

A *density matrix* describing a state of  $X$  is matrix with complex number entries whose rows and columns have been placed in correspondence with  $\Sigma$ .

Typical names for density matrices:  $\rho$ ,  $\sigma$ ,  $\xi$ , ...

## Requirements

1. Density matrices have unit trace:  $\text{Tr}(\rho) = 1$
2. Density matrices are positive semidefinite:  $\rho \geq 0$

# Definition of density matrices

## Requirements

1. Density matrices have unit trace:  $\text{Tr}(\rho) = 1$
2. Density matrices are positive semidefinite:  $\rho \geq 0$

The first condition refers to the **trace**, which is defined for all square matrices as the sum of the diagonal entries.

$$\text{Tr} \begin{pmatrix} \alpha_{0,0} & \alpha_{0,1} & \cdots & \alpha_{0,n-1} \\ \alpha_{1,0} & \alpha_{1,1} & \cdots & \alpha_{1,n-1} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{n-1,0} & \alpha_{n-1,1} & \cdots & \alpha_{n-1,n-1} \end{pmatrix} = \alpha_{0,0} + \alpha_{1,1} + \cdots + \alpha_{n-1,n-1}$$

The trace is a linear function:

$$\text{Tr}(\alpha A + \beta B) = \alpha \text{Tr}(A) + \beta \text{Tr}(B)$$

# Definition of density matrices

## Requirements

1. Density matrices have unit trace:  $\text{Tr}(\rho) = 1$
2. Density matrices are positive semidefinite:  $\rho \geq 0$

The second requirement refers to  $\rho$  being *positive semidefinite*. This is a property that can be expressed in several different ways, including these:

- There exists a matrix  $M$  such that  $\rho = M^\dagger M$ .
- The matrix  $\rho$  is Hermitian, meaning that  $\rho = \rho^\dagger$ , and all of its eigenvalues are nonnegative real numbers.
- For every complex vector  $|\psi\rangle$  we have  $\langle\psi|\rho|\psi\rangle \geq 0$ .

Note that the notation  $\rho \geq 0$  means that  $\rho$  is positive semidefinite — not that each entry of  $\rho$  is nonnegative.

# Examples

## Requirements

1. Density matrices have unit trace:  $\text{Tr}(\rho) = 1$
2. Density matrices are positive semidefinite:  $\rho \geq 0$

## Example of a positive semidefinite matrix

We can generate an example of a positive semidefinite matrix by first choosing a matrix  $M$  arbitrarily and computing  $M^\dagger M$ .

$$M = \begin{pmatrix} -4 - 9i & 8 & 2 + 9i \\ -7 & -4 - 9i & -9 + 7i \\ 1 - 5i & 8 - 6i & -4i \end{pmatrix}$$

$$M^\dagger M = \begin{pmatrix} 172 & 34 + 169i & -6 - 71i \\ 34 - 169i & 261 & 13 - 69i \\ -6 + 71i & 13 + 69i & 231 \end{pmatrix}$$

# Examples

## Requirements

1. Density matrices have unit trace:  $\text{Tr}(\rho) = 1$
2. Density matrices are positive semidefinite:  $\rho \geq 0$

## Examples of density matrices

$$\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$\begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix}$$

$$\begin{pmatrix} \frac{3}{4} & \frac{i}{8} \\ -\frac{i}{8} & \frac{1}{4} \end{pmatrix}$$

$$\begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{pmatrix}$$

$$\frac{1}{664} \begin{pmatrix} 172 & 34 + 169i & -6 - 71i \\ 34 - 169i & 261 & 13 - 69i \\ -6 + 71i & 13 + 69i & 231 \end{pmatrix}$$

# Interpretation

## Requirements

1. Density matrices have unit trace:  $\text{Tr}(\rho) = 1$
2. Density matrices are positive semidefinite:  $\rho \geq 0$

## Intuitive meaning of density matrix entries

For a given density matrix both the rows and the columns correspond to classical states.

- *Diagonal* entries are the probabilities for each classical state to appear from a standard basis measurement.
- *Off-diagonal* entries describe how the two corresponding classical states are *in quantum superposition*.

# Connection to state vectors

A quantum state vector  $|\psi\rangle$  is a column vector having Euclidean norm 1.

Here's the density matrix representation of the same state:

$$|\psi\rangle\langle\psi|$$

States that are represented by density matrices of this form are called *pure states*.

## Examples

$$|+\mathbf{i}\rangle = \frac{1}{\sqrt{2}}|0\rangle + \frac{\mathbf{i}}{\sqrt{2}}|1\rangle = \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{\mathbf{i}}{\sqrt{2}} \end{pmatrix}$$

$$|+\mathbf{i}\rangle\langle+\mathbf{i}| = \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{\mathbf{i}}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{\mathbf{i}}{\sqrt{2}} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & -\frac{\mathbf{i}}{2} \\ \frac{\mathbf{i}}{2} & \frac{1}{2} \end{pmatrix}$$

# Connection to state vectors

A quantum state vector  $|\psi\rangle$  is a column vector having Euclidean norm 1.

Here's the density matrix representation of the same state:

$$|\psi\rangle\langle\psi|$$

States that are represented by density matrices of this form are called *pure states*.

## Examples

$$\begin{aligned} |0\rangle\langle 0| &= \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} & |+\rangle\langle +| &= \begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix} \\ |1\rangle\langle 1| &= \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} & |-\rangle\langle -| &= \begin{pmatrix} \frac{1}{2} & -\frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} \end{pmatrix} \end{aligned}$$

# Connection to state vectors

A quantum state vector  $|\psi\rangle$  is a column vector having Euclidean norm 1.

Here's the density matrix representation of the same state:

$$|\psi\rangle\langle\psi|$$

States that are represented by density matrices of this form are called *pure states*.

## Examples

$$|v\rangle = \frac{1+2i}{3} |0\rangle - \frac{2}{3} |1\rangle$$

$$|v\rangle\langle v| = \begin{pmatrix} \frac{5}{9} & \frac{-2-4i}{9} \\ \frac{-2+4i}{9} & \frac{4}{9} \end{pmatrix}$$

# Connection to state vectors

A quantum state vector  $|\psi\rangle$  is a column vector having Euclidean norm 1.

Here's the density matrix representation of the same state:

$$|\psi\rangle\langle\psi|$$

States that are represented by density matrices of this form are called *pure states*.

In general

$$|\psi\rangle = \begin{pmatrix} \alpha_0 \\ \alpha_1 \\ \vdots \\ \alpha_{n-1} \end{pmatrix} \Rightarrow |\psi\rangle\langle\psi| = \begin{pmatrix} |\alpha_0|^2 & \alpha_0 \overline{\alpha_1} & \cdots & \alpha_0 \overline{\alpha_{n-1}} \\ \alpha_1 \overline{\alpha_0} & |\alpha_1|^2 & \cdots & \alpha_1 \overline{\alpha_{n-1}} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{n-1} \overline{\alpha_0} & \alpha_{n-1} \overline{\alpha_1} & \cdots & |\alpha_{n-1}|^2 \end{pmatrix}$$

# Connection to state vectors

A quantum state vector  $|\psi\rangle$  is a column vector having Euclidean norm 1.

Here's the density matrix representation of the same state:

$$|\psi\rangle\langle\psi|$$

States that are represented by density matrices of this form are called *pure states*.

## Remark

There is no *global phase* degeneracy for density matrices: two quantum states are identical if and only if their density matrix representations are equal.

Suppose  $|\psi\rangle$  and  $|\phi\rangle$  are quantum state vectors that differ by a global phase:

$$|\phi\rangle = e^{i\theta} |\psi\rangle$$

The corresponding density matrices are identical:

$$|\phi\rangle\langle\phi| = (e^{i\theta} |\psi\rangle)(e^{i\theta} |\psi\rangle)^\dagger = e^{i(\theta-\theta)} |\psi\rangle\langle\psi| = |\psi\rangle\langle\psi|$$

# Probabilistic selections

## Key property of density matrices

*Convex combinations* of density matrices represent probabilistic selections of quantum states.

Let  $\rho$  and  $\sigma$  be density matrices representing quantum states of a system, and suppose we prepare the system in state  $\rho$  with probability  $p \in [0, 1]$  and  $\sigma$  with probability  $1 - p$ .

The resulting state is represented by this density matrix:

$$p\rho + (1 - p)\sigma$$

More generally, let  $\rho_0, \dots, \rho_{m-1}$  be density matrices, let  $(p_0, \dots, p_{m-1})$  be a probability vector, and suppose we prepare a system in state  $\rho_k$  with probability  $p_k$ .

The resulting state is represented by this density matrix:

$$\sum_{k=0}^{m-1} p_k \rho_k$$

# Probabilistic selections

## Key property of density matrices

*Convex combinations* of density matrices represent probabilistic selections of quantum states.

More generally, let  $\rho_0, \dots, \rho_{m-1}$  be density matrices, let  $(p_0, \dots, p_{m-1})$  be a probability vector, and suppose we prepare a system in state  $\rho_k$  with probability  $p_k$ .

The resulting state is represented by this density matrix:

$$\sum_{k=0}^{m-1} p_k \rho_k$$

If  $|\psi_0\rangle, \dots, |\psi_{m-1}\rangle$  are quantum state vectors and we prepare a system in state  $|\psi_k\rangle$  with probability  $p_k$ , then the resulting state is represented by this density matrix:

$$\sum_{k=0}^{m-1} p_k |\psi_k\rangle\langle\psi_k|$$

# Probabilistic selections

## Key property of density matrices

*Convex combinations* of density matrices represent probabilistic selections of quantum states.

## Example

A qubit is prepared in the state  $|0\rangle$  with probability  $1/2$  and in the state  $|+\rangle$  with probability  $1/2$ .

Here's the density matrix representation of the resulting state:

$$\frac{1}{2}|0\rangle\langle 0| + \frac{1}{2}|+\rangle\langle +| = \frac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \frac{1}{2}\begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix} = \begin{pmatrix} \frac{3}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} \end{pmatrix}$$

# Completely mixed state

Suppose we set the state of a qubit to be  $|0\rangle$  or  $|1\rangle$  randomly, each with probability  $1/2$ . Here's the density matrix representation of its state.

$$\frac{1}{2}|0\rangle\langle 0| + \frac{1}{2}|1\rangle\langle 1| = \frac{1}{2}\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \frac{1}{2}\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{pmatrix} = \frac{\mathbb{1}}{2}$$

This is known as the *completely mixed state* — it represents complete uncertainty about the state of a qubit.

Now suppose that we change the procedure: in place of the states  $|0\rangle$  and  $|1\rangle$  we'll use the states  $|+\rangle$  and  $|-\rangle$ .

$$\frac{1}{2}|+\rangle\langle +| + \frac{1}{2}|-\rangle\langle -| = \frac{1}{2}\begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix} + \frac{1}{2}\begin{pmatrix} \frac{1}{2} & -\frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{pmatrix} = \frac{\mathbb{1}}{2}$$

It's the same state as before, the completely mixed state. This is a feature, not a bug! The two procedures can't be distinguished by measuring the qubit.

# Probabilistic states

Classical states can be represented by density matrices.

## Example

Density matrices for the classical states 0 and 1 of a qubit:

$$|0\rangle\langle 0| = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad |1\rangle\langle 1| = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

## In general

If a system has classical state set  $\Sigma$ , then each classical state  $\alpha \in \Sigma$  is represented by the density matrix  $|\alpha\rangle\langle \alpha|$ .

# Probabilistic states

## In general

If a system has classical state set  $\Sigma$ , then each classical state  $\alpha \in \Sigma$  is represented by the density matrix  $|\alpha\rangle\langle\alpha|$ .

Probabilistic states are represented by **convex combinations** of such density matrices.

## Example

Suppose the classical state set of  $X$  is  $\{0, \dots, n-1\}$ .

We can identify the probabilistic state of  $X$  represented by the probability vector  $(p_0, \dots, p_{n-1})$  with this density matrix:

$$\rho = \sum_{k=0}^{n-1} p_k |k\rangle\langle k| = \begin{pmatrix} p_0 & 0 & \cdots & 0 \\ 0 & p_1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & p_{n-1} \end{pmatrix}$$

# Spectral theorem

## Spectral theorem for positive semidefinite matrices

Suppose  $P$  is an  $n \times n$  positive semidefinite matrix.

There exists an orthonormal basis  $\{|\psi_0\rangle, \dots, |\psi_{n-1}\rangle\}$  of vectors along with *nonnegative real* numbers  $\lambda_0, \dots, \lambda_{n-1}$  such that

$$P = \sum_{k=0}^{n-1} \lambda_k |\psi_k\rangle\langle\psi_k|$$

## Implication for density matrices

Any  $n \times n$  density matrix  $\rho$  can be expressed as

$$\rho = \sum_{k=0}^{n-1} p_k |\psi_k\rangle\langle\psi_k|$$

for an orthonormal basis  $\{|\psi_0\rangle, \dots, |\psi_{n-1}\rangle\}$  and a probability vector  $(p_0, \dots, p_{n-1})$ .

# Spectral theorem

## Example

Recall the example where a qubit is prepared in the state  $|0\rangle$  with probability  $1/2$  and  $|+\rangle$  with probability  $1/2$ .

$$\frac{1}{2}|0\rangle\langle 0| + \frac{1}{2}|+\rangle\langle +| = \begin{pmatrix} \frac{3}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} \end{pmatrix}$$

This density matrix can alternatively be expressed as follows:

$$\begin{pmatrix} \frac{3}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} \end{pmatrix} = \cos^2\left(\frac{\pi}{8}\right)|\psi_{\pi/8}\rangle\langle\psi_{\pi/8}| + \sin^2\left(\frac{\pi}{8}\right)|\psi_{5\pi/8}\rangle\langle\psi_{5\pi/8}|$$

$$|\psi_{\alpha}\rangle = \cos(\alpha)|0\rangle + \sin(\alpha)|1\rangle$$

# Qubit quantum state vectors

Up to a *global phase*, every qubit quantum state vector is equivalent to

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right) |0\rangle + e^{i\phi} \sin\left(\frac{\theta}{2}\right) |1\rangle$$

for  $\theta \in [0, \pi]$  and  $\phi \in [0, 2\pi)$ .

## Almost uniqueness

- If  $\theta = 0$  or  $\theta = \pi$ , then  $\phi$  is *irrelevant*.
- If  $\theta \in (0, \pi)$ , then  $\phi$  is *unique*.

# Pure states of a qubit

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + e^{i\phi}\sin\left(\frac{\theta}{2}\right)|1\rangle \quad \theta \in [0, \pi] \quad \phi \in [0, 2\pi)$$

Here's this state's density matrix representation:

$$|\psi\rangle\langle\psi| = \begin{pmatrix} \cos^2\left(\frac{\theta}{2}\right) & e^{-i\phi}\cos\left(\frac{\theta}{2}\right)\sin\left(\frac{\theta}{2}\right) \\ e^{i\phi}\cos\left(\frac{\theta}{2}\right)\sin\left(\frac{\theta}{2}\right) & \sin^2\left(\frac{\theta}{2}\right) \end{pmatrix}$$

Like all  $2 \times 2$  matrices, there's a unique way to express this density matrix as a *linear combination of Pauli matrices*.

$$\mathbb{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad \sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$
$$|\psi\rangle\langle\psi| = \frac{\mathbb{1} + \sin(\theta)\cos(\phi)\sigma_x + \sin(\theta)\sin(\phi)\sigma_y + \cos(\theta)\sigma_z}{2}$$

$$\cos^2\left(\frac{\theta}{2}\right) = \frac{1 + \cos(\theta)}{2}$$

$$\sin^2\left(\frac{\theta}{2}\right) = \frac{1 - \cos(\theta)}{2}$$


$$\cos\left(\frac{\theta}{2}\right)\sin\left(\frac{\theta}{2}\right) = \frac{\sin(\theta)}{2}$$

$$e^{i\phi} = \cos(\phi) + i\sin(\phi)$$

# Bloch sphere

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + e^{i\phi}\sin\left(\frac{\theta}{2}\right)|1\rangle \quad \theta \in [0, \pi] \quad \phi \in [0, 2\pi)$$


$$|\psi\rangle\langle\psi| = \frac{\mathbb{1} + \sin(\theta)\cos(\phi)\sigma_x + \sin(\theta)\sin(\phi)\sigma_y + \cos(\theta)\sigma_z}{2}$$



# Bloch sphere

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + e^{i\phi}\sin\left(\frac{\theta}{2}\right)|1\rangle \quad \theta \in [0, \pi] \quad \phi \in [0, 2\pi)$$

$$|\psi\rangle\langle\psi| = \frac{\mathbb{1} + \sin(\theta)\cos(\phi)\sigma_x + \sin(\theta)\sin(\phi)\sigma_y + \cos(\theta)\sigma_z}{2}$$




This is the *Bloch sphere* representation of the qubit state  $|\psi\rangle$ .

# Bloch sphere examples

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + e^{i\phi}\sin\left(\frac{\theta}{2}\right)|1\rangle \quad \theta \in [0, \pi] \quad \phi \in [0, 2\pi)$$

$$|\psi\rangle\langle\psi| = \frac{\mathbb{1} + \sin(\theta)\cos(\phi)\sigma_x + \sin(\theta)\sin(\phi)\sigma_y + \cos(\theta)\sigma_z}{2}$$



## Examples

The standard basis  $\{|0\rangle, |1\rangle\}$

$$|0\rangle = \cos(0)|0\rangle + e^{i\phi}\sin(0)|1\rangle$$

$$|1\rangle = \cos\left(\frac{\pi}{2}\right)|0\rangle + e^{i\phi}\sin\left(\frac{\pi}{2}\right)|1\rangle$$


$$|0\rangle\langle 0| = \frac{\mathbb{1} + \sigma_z}{2}$$

$$|1\rangle\langle 1| = \frac{\mathbb{1} - \sigma_z}{2}$$

# Bloch sphere examples

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + e^{i\phi}\sin\left(\frac{\theta}{2}\right)|1\rangle \quad \theta \in [0, \pi] \quad \phi \in [0, 2\pi)$$

$$|\psi\rangle\langle\psi| = \frac{\mathbb{1} + \sin(\theta)\cos(\phi)\sigma_x + \sin(\theta)\sin(\phi)\sigma_y + \cos(\theta)\sigma_z}{2}$$



## Examples

The basis  $\{|+\rangle, |-\rangle\}$

$$|+\rangle = \cos\left(\frac{\pi}{4}\right)|0\rangle + e^{i0}\sin\left(\frac{\pi}{4}\right)|1\rangle$$

$$|-\rangle = \cos\left(\frac{\pi}{4}\right)|0\rangle + e^{i\pi}\sin\left(\frac{\pi}{4}\right)|1\rangle$$


$$|+\rangle\langle+| = \frac{\mathbb{1} + \sigma_x}{2}$$

$$|-\rangle\langle-| = \frac{\mathbb{1} - \sigma_x}{2}$$

# Bloch sphere examples

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + e^{i\phi}\sin\left(\frac{\theta}{2}\right)|1\rangle \quad \theta \in [0, \pi] \quad \phi \in [0, 2\pi)$$

$$|\psi\rangle\langle\psi| = \frac{\mathbb{1} + \sin(\theta)\cos(\phi)\sigma_x + \sin(\theta)\sin(\phi)\sigma_y + \cos(\theta)\sigma_z}{2}$$



## Examples

The basis  $\{|+i\rangle, |-i\rangle\}$

$$|+i\rangle = \frac{1}{\sqrt{2}}|0\rangle + \frac{i}{\sqrt{2}}|1\rangle$$

$$|-i\rangle = \frac{1}{\sqrt{2}}|0\rangle - \frac{i}{\sqrt{2}}|1\rangle$$


$$|+i\rangle = \cos\left(\frac{\pi}{4}\right)|0\rangle + e^{i\frac{\pi}{2}}\sin\left(\frac{\pi}{4}\right)|1\rangle$$

$$|-i\rangle = \cos\left(\frac{\pi}{4}\right)|0\rangle + e^{i\frac{3\pi}{2}}\sin\left(\frac{\pi}{4}\right)|1\rangle$$

# Bloch sphere examples

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + e^{i\phi}\sin\left(\frac{\theta}{2}\right)|1\rangle \quad \theta \in [0, \pi] \quad \phi \in [0, 2\pi)$$

$$|\psi\rangle\langle\psi| = \frac{\mathbb{1} + \sin(\theta)\cos(\phi)\sigma_x + \sin(\theta)\sin(\phi)\sigma_y + \cos(\theta)\sigma_z}{2}$$



## Examples

The basis  $\{|+i\rangle, |-i\rangle\}$

$$|+i\rangle = \cos\left(\frac{\pi}{4}\right)|0\rangle + e^{i\frac{\pi}{2}}\sin\left(\frac{\pi}{4}\right)|1\rangle$$

$$|-i\rangle = \cos\left(\frac{\pi}{4}\right)|0\rangle + e^{i\frac{3\pi}{2}}\sin\left(\frac{\pi}{4}\right)|1\rangle$$


$$|+i\rangle\langle+i| = \frac{\mathbb{1} + \sigma_y}{2}$$

$$|-i\rangle\langle-i| = \frac{\mathbb{1} - \sigma_y}{2}$$

# Bloch sphere examples

$$|\psi\rangle = \cos\left(\frac{\theta}{2}\right)|0\rangle + e^{i\phi}\sin\left(\frac{\theta}{2}\right)|1\rangle \quad \theta \in [0, \pi] \quad \phi \in [0, 2\pi)$$

$$|\psi\rangle\langle\psi| = \frac{\mathbb{1} + \sin(\theta)\cos(\phi)\sigma_x + \sin(\theta)\sin(\phi)\sigma_y + \cos(\theta)\sigma_z}{2}$$



## Examples

Another class of state vectors:


$$|\psi_\alpha\rangle = \cos(\alpha)|0\rangle + \sin(\alpha)|1\rangle, \quad \alpha \in [0, \pi)$$

$$|\psi_\alpha\rangle\langle\psi_\alpha| = \frac{\mathbb{1} + \sin(2\alpha)\sigma_x + \cos(2\alpha)\sigma_z}{2}$$

# Bloch ball

The **Bloch ball** contains all of the points on the Bloch sphere as well as those inside it.

- Points in the interior of the Bloch ball correspond to qubit states that are not pure.
- Convex combinations of points in the Bloch ball correspond to convex combinations of qubit density matrices.



## Examples


The completely mixed state lies at the center of the Bloch ball.

$$\begin{aligned}\frac{\mathbb{1}}{2} &= \frac{1}{2}|0\rangle\langle 0| + \frac{1}{2}|1\rangle\langle 1| \\ &= \frac{1}{2}|+\rangle\langle +| + \frac{1}{2}|-\rangle\langle -| \\ &= \frac{1}{2}|+i\rangle\langle +i| + \frac{1}{2}|-i\rangle\langle -i|\end{aligned}$$

# Bloch ball

The **Bloch ball** contains all of the points on the Bloch sphere as well as those inside it.

- Points in the interior of the Bloch ball correspond to qubit states that are not pure.
- Convex combinations of points in the Bloch ball correspond to convex combinations of qubit density matrices.



## Examples


The completely mixed state lies at the center of the Bloch ball.

$$\frac{1}{2} = \frac{1 + 0 \cdot \sigma_x + 0 \cdot \sigma_y + 0 \cdot \sigma_z}{2}$$

# Bloch ball

The **Bloch ball** contains all of the points on the Bloch sphere as well as those inside it.

- Points in the interior of the Bloch ball correspond to qubit states that are not pure.
- Convex combinations of points in the Bloch ball correspond to convex combinations of qubit density matrices.



## Examples

$$\begin{pmatrix} \frac{3}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} \end{pmatrix} = \frac{1}{2}|0\rangle\langle 0| + \frac{1}{2}|+\rangle\langle +|$$

$$= \cos^2\left(\frac{\pi}{8}\right)|\psi_{\pi/8}\rangle\langle\psi_{\pi/8}|$$

$$+ \sin^2\left(\frac{\pi}{8}\right)|\psi_{5\pi/8}\rangle\langle\psi_{5\pi/8}|$$

# Multiple systems

Density matrices can represent states of multiple systems:

- Multiple system are viewed as single, compound systems.
- The rows and columns of density matrices for multiple systems correspond to *Cartesian products* of the classical state sets of the individual systems.

Example: Bell states

$$|\Phi^+\rangle\langle\Phi^+| = \begin{pmatrix} \frac{1}{2} & 0 & 0 & \frac{1}{2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \frac{1}{2} & 0 & 0 & \frac{1}{2} \end{pmatrix}$$

$$|\Phi^-\rangle\langle\Phi^-| = \begin{pmatrix} \frac{1}{2} & 0 & 0 & -\frac{1}{2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -\frac{1}{2} & 0 & 0 & \frac{1}{2} \end{pmatrix}$$

$$|\Psi^+\rangle\langle\Psi^+| = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$|\Psi^-\rangle\langle\Psi^-| = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & -\frac{1}{2} & 0 \\ 0 & -\frac{1}{2} & \frac{1}{2} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

# Independence and correlation

**Tensor products** of density matrices represent **independence** between systems.

## Product states

If X is prepared in the state  $\rho$  and Y is independently prepared in the state  $\sigma$ , then the state of (X, Y) is the tensor product  $\rho \otimes \sigma$ .

States of this form are called **product states**.

Density matrices that cannot be expressed as product states represent **correlations** between systems. Different types of correlations can be considered.

## Example: a correlated classical state

A uniform random bit shared between Alice and Bob has this state.

$$\frac{1}{2}|0\rangle\langle 0| \otimes |0\rangle\langle 0| + \frac{1}{2}|1\rangle\langle 1| \otimes |1\rangle\langle 1| = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2} \end{pmatrix}$$

# Independence and correlation

**Tensor products** of density matrices represent **independence** between systems.

## Product states

If  $X$  is prepared in the state  $\rho$  and  $Y$  is independently prepared in the state  $\sigma$ , then the state of  $(X, Y)$  is the tensor product  $\rho \otimes \sigma$ .

States of this form are called **product states**.

Density matrices that cannot be expressed as product states represent **correlations** between systems. Different types of correlations can be considered.

## Example: ensembles of quantum states

An **ensemble** describes a random selection of a quantum state.

The following density matrix represents an ensemble, assuming  $(p_0, \dots, p_{m-1})$  is a probability vector and  $\rho_0, \dots, \rho_{m-1}$  are density matrices of the same size.

$$\sum_{k=0}^{m-1} p_k |k\rangle\langle k| \otimes \rho_k$$

# Independence and correlation

**Tensor products** of density matrices represent **independence** between systems.

## Product states

If X is prepared in the state  $\rho$  and Y is independently prepared in the state  $\sigma$ , then the state of (X, Y) is the tensor product  $\rho \otimes \sigma$ .

States of this form are called **product states**.

Density matrices that cannot be expressed as product states represent **correlations** between systems. Different types of correlations can be considered.

## Example: separable states

Convex combinations of product states represent classical correlations among quantum states.

$$\sum_{k=0}^{m-1} p_k \rho_k \otimes \sigma_k$$

States that can be written like this are **separable**; all the rest are **entangled**.

# Reduced states for an e-bit

Alice and Bob *share an e-bit*: Alice holds A, Bob holds B, and (A, B) is in this state:

$$|\phi^+\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$$

## Question

What is the state of Alice's qubit A in isolation?

Imagine Bob performs a standard basis measurement on B.

Outcome	Probability	Resulting state of A
0	$\frac{1}{2}$	$ 0\rangle$
1	$\frac{1}{2}$	$ 1\rangle$

This leaves the qubit A in the completely mixed state:

$$\frac{1}{2}|0\rangle\langle 0| + \frac{1}{2}|1\rangle\langle 1| = \frac{\mathbb{1}}{2}$$

# Reduced states for an e-bit

Alice and Bob *share an e-bit*: Alice holds A, Bob holds B, and (A, B) is in this state:

$$|\phi^+\rangle = \frac{1}{\sqrt{2}}|00\rangle + \frac{1}{\sqrt{2}}|11\rangle$$

## Question

What is the state of Alice's qubit A in isolation?

## Answer

The completely mixed state  $\frac{1}{2}$ .

# Reduced states in general

Suppose we have a pair of systems (A, B) in a pure state  $|\psi\rangle$ .

Let  $\Gamma$  be the classical state set of B. There is a unique collection  $\{|\phi_b\rangle : b \in \Gamma\}$  of vectors for which this expression is true:

$$|\psi\rangle = \sum_{b \in \Gamma} |\phi_b\rangle \otimes |b\rangle$$

In particular, each  $|\phi_b\rangle$  is given by this formula:

$$|\phi_b\rangle = (\mathbb{1}_A \otimes \langle b|)|\psi\rangle$$

# Reduced states in general

Suppose we have a pair of systems (A, B) in a pure state  $|\psi\rangle$ .

Let  $\Gamma$  be the classical state set of B. There is a unique collection  $\{|\phi_b\rangle : b \in \Gamma\}$  of vectors for which this expression is true:

$$|\psi\rangle = \sum_{b \in \Gamma} |\phi_b\rangle \otimes |b\rangle$$

$$|\phi_b\rangle = (\mathbb{1}_A \otimes \langle b|) |\psi\rangle$$

Suppose a standard basis measurement were performed on B:

- Each  $b \in \Gamma$  appears with probability  $\| |\phi_b\rangle \|^2$ .
- Conditioned on obtaining the outcome  $b$ , the state of A becomes  $\frac{|\phi_b\rangle}{\| |\phi_b\rangle \|}$ .

The reduced state of A:

$$\sum_{b \in \Gamma} \| |\phi_b\rangle \|^2 \frac{|\phi_b\rangle \langle \phi_b|}{\| |\phi_b\rangle \|^2} = \sum_{b \in \Gamma} |\phi_b\rangle \langle \phi_b|$$

# Reduced states in general

Suppose we have a pair of systems (A, B) in a pure state  $|\psi\rangle$ .

Let  $\Gamma$  be the classical state set of B. There is a unique collection  $\{|\phi_b\rangle : b \in \Gamma\}$  of vectors for which this expression is true:

$$|\psi\rangle = \sum_{b \in \Gamma} |\phi_b\rangle \otimes |b\rangle$$

$$|\phi_b\rangle = (\mathbb{1}_A \otimes \langle b|) |\psi\rangle$$

Suppose a standard basis measurement were performed on B:

- Each  $b \in \Gamma$  appears with probability  $\| |\phi_b\rangle \|^2$ .
- Conditioned on obtaining the outcome  $b$ , the state of A becomes  $\frac{|\phi_b\rangle}{\| |\phi_b\rangle \|}$ .

The reduced state of A:

$$\sum_{b \in \Gamma} |\phi_b\rangle \langle \phi_b| = \sum_{b \in \Gamma} (\mathbb{1}_A \otimes \langle b|) |\psi\rangle \langle \psi| (\mathbb{1}_A \otimes |b\rangle)$$

# Reduced states in general

Suppose we have a pair of systems (A, B) in a pure state  $|\psi\rangle$ .

The reduced state of A:

$$\sum_{b \in \Gamma} (\mathbb{1}_A \otimes \langle b |) |\psi\rangle \langle \psi| (\mathbb{1}_A \otimes |b\rangle)$$

Suppose (A, B) is in a state described by a density matrix  $\rho$ . The reduced state of A is described by this density matrix:

$$\rho_A = \sum_{b \in \Gamma} (\mathbb{1}_A \otimes \langle b |) \rho (\mathbb{1}_A \otimes |b\rangle)$$

The reduced state of B is obtained in a similar way.

$$\rho_B = \sum_{a \in \Sigma} (\langle a | \otimes \mathbb{1}_B) \rho (|a\rangle \otimes \mathbb{1}_B)$$

( $\Sigma$  is the classical state set of A.)

# Reduced states in general

Suppose  $(A, B)$  is in a state described by a density matrix  $\rho$ . The reduced state of  $A$  is described by this density matrix:

$$\rho_A = \sum_{b \in \Gamma} (\mathbb{1}_A \otimes \langle b|) \rho (\mathbb{1}_A \otimes |b\rangle)$$

The reduced state of  $B$  is obtained in a similar way.

$$\rho_B = \sum_{a \in \Sigma} (\langle a| \otimes \mathbb{1}_B) \rho (|a\rangle \otimes \mathbb{1}_B)$$

These notions can be generalized to three or more systems in a natural way.

## Example

Suppose  $(A, B, C)$  is in the state  $\rho$ . The reduced states of  $(A, C)$  and  $C$ :

$$\begin{aligned} \rho_{AC} &= \sum_{b \in \Gamma} (\mathbb{1}_A \otimes \langle b| \otimes \mathbb{1}_C) \rho (\mathbb{1}_A \otimes |b\rangle \otimes \mathbb{1}_C) \\ \rho_C &= \sum_{a \in \Sigma} \sum_{b \in \Gamma} (\langle a| \otimes \langle b| \otimes \mathbb{1}_C) \rho (|a\rangle \otimes |b\rangle \otimes \mathbb{1}_C) \end{aligned}$$

# The partial trace

## Definition

$$\text{Tr}_B(\rho) = \sum_{b \in \Gamma} (\mathbb{1}_A \otimes \langle b |) \rho (\mathbb{1}_A \otimes | b \rangle)$$

$$\text{Tr}_A(\rho) = \sum_{a \in \Sigma} (\langle a | \otimes \mathbb{1}_B) \rho (| a \rangle \otimes \mathbb{1}_B)$$

## Equivalent definition

$\text{Tr}_A$  and  $\text{Tr}_B$  are the unique linear mappings for which these equations are always true:

$$\text{Tr}_A(M \otimes N) = \text{Tr}(M)N$$

$$\text{Tr}_B(M \otimes N) = \text{Tr}(N)M$$

## Example

Consider this state of a pair of qubits (A, B):

$$\rho = \frac{1}{2} |0\rangle\langle 0| \otimes |0\rangle\langle 0| + \frac{1}{2} |1\rangle\langle 1| \otimes |+\rangle\langle +|$$

$$\rho_A = \text{Tr}_B(\rho) = \frac{1}{2} |0\rangle\langle 0| + \frac{1}{2} |1\rangle\langle 1| \quad \rho_B = \text{Tr}_A(\rho) = \frac{1}{2} |0\rangle\langle 0| + \frac{1}{2} |+\rangle\langle +|$$